



“Beyond the Blood Pressure Reading: The Pivotal Role of Community Health Nurses in Hypertension Detection, Prevention, Treatment Support, and Long-Term Control”

Alka Shiv¹, Dr. Nitin Chicholkar²

¹PhD Research Scholar, ²Research Supervisor

^{1,2}Index Nursing College,

Malwanchal University, Indore

Date of publication: 15/02/2024

DOI 10.5281/zenodo.22993380

Abstract: Hypertension is one of the most important modifiable risk factors for cardiovascular disease, stroke, chronic kidney disease, heart failure, and premature mortality. Despite the availability of inexpensive diagnostic methods and effective antihypertensive therapies, substantial gaps remain in hypertension awareness, treatment initiation, adherence, follow-up, and sustained blood pressure control. Global evidence indicates that the number of adults living with hypertension has increased markedly, while many affected individuals remain undiagnosed or inadequately treated [1,2]. Because hypertension is frequently asymptomatic, community-level detection and continuous follow-up are essential components of an effective public health response. Community health nurses occupy a strategically important position in this continuum because they can combine clinical assessment, health education, behavioural counselling, family engagement, medication-support activities, home visits, referral, follow-up, and coordination with primary healthcare teams. Evidence from nurse-led interventions demonstrates that structured nursing care can contribute to improvements in blood pressure and cardiovascular risk-factor management [3,4]. Community-based programmes involving trained health workers have also demonstrated improvements in hypertension screening, identification, monitoring, and medication adherence, particularly in resource-constrained settings [5,6]. In India, initiatives such as the India Hypertension Control Initiative demonstrate the value of standardized treatment protocols, team-based care, patient-centred approaches, reliable medication supply, and information systems [7]. This narrative review examines the role of community health nurses in hypertension detection and control, including screening, accurate blood pressure measurement, risk assessment, lifestyle modification, medication adherence, home-based care, referral, digital health, documentation, family participation, and programme monitoring. The review also discusses barriers, professional competencies, ethical considerations, and future directions for strengthening nurse-led hypertension services within primary healthcare.

Keywords: Hypertension; Community Health Nursing; Blood Pressure; Hypertension Screening; Blood Pressure Control; Nurse-Led Intervention; Primary Health Care; Health Education; Medication Adherence; Lifestyle Modification; Cardiovascular Disease Prevention; Community-Based Care.

Introduction

Hypertension is a chronic noncommunicable disease characterized by persistently elevated blood pressure and is a major contributor to cardiovascular morbidity and mortality. It is often described as a “silent” condition because many individuals do not experience recognizable symptoms until complications develop. Consequently, a person may live with elevated blood pressure for years without knowing that

treatment or lifestyle modification is required. The World Health Organization (WHO) reported in 2023 that approximately 1.3 billion adults were living with hypertension and that nearly half of people with hypertension were unaware of their condition. The report emphasized that prevention, early detection, and effective management at the primary-care level are among the most important



opportunities for reducing the consequences of uncontrolled blood pressure [1,2].

The global burden of hypertension is particularly important for low- and middle-income countries, where healthcare systems may experience shortages of trained personnel, limited access to medicines, inadequate continuity of care, and substantial socioeconomic barriers. A large pooled analysis of 1201 population-representative studies involving more than 100 million participants demonstrated substantial differences between countries in hypertension detection, treatment, and control [8]. Globally, among adults aged 30–79 years with hypertension in 2019, only a minority had their blood pressure controlled below 140/90 mmHg [8].

Hypertension management therefore cannot be restricted to occasional clinical encounters. It requires a continuum of care beginning with population awareness and screening and extending through diagnosis, treatment initiation, adherence support, lifestyle modification, monitoring, and long-term follow-up. Community health nurses are particularly suited to this continuum because they work close to individuals, families, schools, workplaces, community organizations, and primary healthcare facilities. Their accessibility provides opportunities to identify people who rarely attend hospitals or clinics and to support those who have already been diagnosed but experience difficulties maintaining treatment. The contemporary approach to hypertension control emphasizes standardized treatment protocols, team-based care, patient-centred services, access to essential medicines and technologies, and information systems for monitoring [9]. Within such models, the community health nurse can function not merely as a provider of health education but as an active participant in case identification, risk assessment, care coordination, treatment support, behavioural counselling, follow-up, and quality improvement.

Hypertension as a Community Health Priority

The importance of community-based hypertension control arises from the relationship between elevated blood pressure and major cardiovascular and renal outcomes. Persistent hypertension contributes to vascular injury and increases the risk of stroke, myocardial infarction, heart failure, chronic

kidney disease, peripheral vascular disease, and other complications [1,2]. The long-term consequences are not limited to individual health. Families may experience financial strain because of treatment expenses, hospitalization, disability, loss of income, and caregiving responsibilities.

Hypertension is also strongly associated with modifiable behavioural and environmental factors. High dietary sodium intake, unhealthy dietary patterns, inadequate physical activity, tobacco use, excessive alcohol consumption, obesity, and psychosocial stress can contribute to increased cardiovascular risk. The WHO HEARTS framework specifically incorporates healthy-lifestyle counselling as one of the major components of comprehensive cardiovascular disease prevention and management [9,10]. Community health nurses can address these factors repeatedly and in context, rather than relying on one-time counselling in a hospital setting.

The community setting is particularly important because risk factors are embedded within everyday life. Dietary counselling, for example, becomes more meaningful when the nurse understands local foods, cooking practices, household decision-making, food purchasing patterns, cultural preferences, and economic limitations. Similarly, advice concerning physical activity should consider age, occupation, safety, disability, environmental conditions, and available community resources. Such contextualized health education can make hypertension prevention and control more practical.

Community Health Nurses and Early Detection of Hypertension

Early detection represents one of the most important contributions of community health nursing. Since hypertension can remain asymptomatic, blood pressure measurement provides a practical means of identifying individuals who may require further assessment. Community health nurses can incorporate blood pressure measurement into home visits, community outreach programmes, health camps, primary healthcare clinics, workplace programmes, older-adult services, and other community encounters.



Screening should be performed using validated equipment and standardized procedures. The individual should be appropriately positioned, allowed to rest before measurement, and assessed using an appropriately sized cuff. Multiple readings may be necessary because a single elevated reading does not always establish a diagnosis. Community health nurses should recognize factors that can temporarily influence blood pressure, including recent physical activity, emotional stress, caffeine intake, tobacco use, pain, and an improperly performed measurement. In India, an expert consensus statement recommended blood pressure screening beginning at adulthood and emphasized appropriate equipment and repeated measurements during screening [11].

The nurse's role extends beyond recording a numerical value. A person with an elevated reading should receive an appropriate explanation and, according to the applicable clinical protocol, repeat measurement, further assessment, or referral. Nurses should avoid labelling an individual as hypertensive solely on the basis of one abnormal screening value unless the relevant protocol specifically permits such classification. Accurate documentation and appropriate referral are essential components of safe screening.

Community-based screening can also reduce inequalities in access to diagnosis. People who are older, economically disadvantaged, geographically isolated, socially marginalized, or reluctant to visit healthcare facilities may be missed by facility-based services. Community nurses can bring screening closer to such populations.

Accurate Blood Pressure Measurement as a Nursing Responsibility

Accurate blood pressure measurement is fundamental to hypertension detection and control. An incorrect reading can lead to unnecessary anxiety, inappropriate treatment, or failure to recognize genuine hypertension. Community health nurses therefore require competency in equipment selection, cuff placement, patient positioning, repeated measurement, documentation, and interpretation within the appropriate clinical framework.

The nurse should ensure that the person is seated comfortably, with appropriate support, and that the arm is positioned appropriately. The correct cuff size should be selected according to the individual's arm circumference. Where indicated by protocol, measurements should be obtained on more than one occasion or repeated after an appropriate interval. Equipment should be maintained and periodically checked according to institutional requirements. Community nurses should also educate individuals about home blood pressure monitoring when it is appropriate. Home monitoring can provide additional information about blood pressure patterns and may support patient participation in self-management. However, patients should be taught how to use the device correctly and how to record readings. Community nurses can review these records during home visits or clinic appointments and communicate concerning findings to the appropriate healthcare professional.

Risk Assessment and Identification of Vulnerable Individuals

Hypertension screening should be integrated with broader cardiovascular risk assessment rather than viewed as an isolated activity. During community encounters, nurses can assess important risk factors such as age, family history, obesity, diabetes, dyslipidaemia, tobacco use, physical inactivity, dietary patterns, alcohol consumption, previous cardiovascular disease, kidney disease, and medication history.

A family-centred assessment is valuable because lifestyle practices frequently occur at the household level. For example, dietary sodium reduction may be difficult if the household prepares high-salt foods for everyone. Involving family members can therefore increase the feasibility of lifestyle modification.

Community health nurses should also identify individuals who may require priority referral. Markedly elevated blood pressure, symptoms suggestive of acute cardiovascular or neurological complications, pregnancy-associated hypertension, suspected secondary hypertension, medication-related adverse effects, or persistent uncontrolled blood pressure despite treatment require



appropriate clinical evaluation. The nurse should understand the limits of independent practice and promptly refer patients according to local protocols.

Health Education and Lifestyle Modification

Health education is one of the most visible and potentially influential components of community health nursing. However, effective education should move beyond simply telling patients what they should do. The nurse should assess existing knowledge, identify barriers, establish realistic goals, and provide repeated reinforcement.

Dietary modification is an important area of counselling. Nurses can encourage reduced sodium intake, greater consumption of fruits and vegetables where appropriate, balanced dietary patterns, moderation of processed foods, and healthier food preparation practices. Rather than presenting complex dietary prescriptions, community nurses can translate recommendations into practical household behaviours.

Physical activity counselling should be individualized. Adults without contraindications can be encouraged to engage in regular physical activity appropriate to their health status. Nurses can promote walking groups, active transportation, household activity, community exercise programmes, and gradual increases in daily movement. For older adults or people with chronic conditions, the nurse should encourage activity consistent with professional assessment and safety considerations.

Tobacco cessation and reduction of harmful alcohol consumption should form part of cardiovascular risk reduction. Nurses can provide brief interventions, assess readiness to change, reinforce motivation, and facilitate referral to appropriate cessation services. Weight management should be approached sensitively and should focus on sustainable behavioural changes rather than stigmatizing language.

The WHO HEARTS healthy-lifestyle counselling module emphasizes brief interventions addressing major cardiovascular risk behaviours [10]. Community nurses can integrate these interventions into routine encounters instead of treating lifestyle counselling as a separate activity.

Medication Adherence and Treatment Support

Effective antihypertensive medication is available for most individuals who require pharmacological treatment, but treatment effectiveness depends substantially on consistent use and appropriate follow-up. Community health nurses can identify practical and behavioural reasons for non-adherence. These may include forgetfulness, complicated medication schedules, cost, side effects, lack of understanding, misconceptions, fear of lifelong treatment, perceived absence of symptoms, transportation problems, medicine shortages, or competing family responsibilities.

The nurse should provide clear explanations about the purpose of prescribed medication and the importance of continuing treatment even when the individual feels well. Patients should be encouraged not to discontinue or modify prescribed medication independently. When adverse effects or other concerns are reported, the nurse should communicate them to the prescribing clinician or follow the relevant treatment protocol.

WHO's 2021 guideline provides recommendations concerning pharmacological treatment, follow-up intervals, blood pressure targets, and treatment approaches for adults with hypertension [12]. Community nurses should apply these recommendations through locally authorized protocols rather than independently prescribing or changing medication outside their scope of practice.

Home Visits and Continuity of Care

Home visiting provides community health nurses with an opportunity to understand the circumstances affecting hypertension control. During a home visit, the nurse can assess medication organization, family support, diet, physical activity, smoking or alcohol-related behaviours, access to healthcare, and barriers to follow-up. Home-based care can be particularly valuable for older adults, people with mobility limitations, individuals living alone, and patients who repeatedly miss clinic appointments.

Continuity is an important advantage of community nursing. Hypertension is not a condition that can be effectively managed through a single educational session. Repeated



contacts allow nurses to assess progress, reinforce positive changes, identify emerging problems, and coordinate referrals. A community nurse may therefore function as a consistent point of contact within the broader primary healthcare team.

Evidence from community-based nursing interventions suggests that nurse-led approaches can improve self-care behaviours and patient satisfaction, although effects on blood pressure may vary according to intervention design and duration [13].

Evidence Supporting Nurse-Led Hypertension Management

The evidence base for nurse-led hypertension care has expanded considerably. An early systematic review and meta-analysis found that nurse-led interventions incorporating structured treatment algorithms, prescribing where legally authorized, telephone monitoring, and community monitoring were associated with improvements in blood pressure outcomes compared with usual care [3].

More recent evidence has similarly demonstrated potential benefits. A 2022 systematic review and meta-analysis of 11 randomized trials involving 4454 participants reported significant reductions in systolic and diastolic blood pressure in several nurse-led interventions and improvements in medication adherence and lifestyle-related outcomes [4]. These findings support the concept that nurses can make a meaningful contribution when their responsibilities are clearly structured and integrated into primary care.

Digital approaches provide another emerging opportunity. A systematic review of nurse-led digital health interventions found reductions in systolic and diastolic blood pressure compared with usual care, together with improvements in self-management and medication adherence [14]. Telephonic follow-up, digital reminders, electronic records, remote blood pressure reporting, and patient education platforms can extend the reach of community nursing.

The evidence should nevertheless be interpreted carefully because nurse-led interventions differ substantially in intensity, professional authority, duration, clinical context, and combination of educational, behavioural, and

medication-related components. The evidence supports nursing involvement as part of team-based hypertension management rather than suggesting that one model is universally applicable.

Community Health Workers and the Extended Nursing Team

Community health nurses frequently work with community health workers and other frontline personnel. Task sharing can expand hypertension detection and follow-up, particularly where professional nursing resources are limited. WHO's HEARTS approach explicitly recognizes team-based care and task sharing as important elements of primary-care hypertension programmes [9,15].

A systematic review of community health worker interventions in low- and middle-income countries found that interventions commonly included blood pressure measurement, health education, lifestyle support, medication adherence assistance, referral, and follow-up [5]. Such programmes can be strengthened when nurses provide training, supervision, clinical escalation pathways, and quality assurance.

Indian evidence provides a particularly relevant example. In Chhattisgarh, an intervention that equipped community health workers to measure blood pressure was associated with substantially greater screening, identification, recent blood pressure measurement, and medication adherence compared with the comparison group [6]. This illustrates how community-level personnel can extend the reach of hypertension services when appropriately trained and connected with the formal healthcare system.

Role of Community Health Nurses in India

India has a substantial hypertension burden and has developed large-scale strategies for improving detection and control. The India Hypertension Control Initiative (IHCI) incorporates standardized treatment protocols, reliable availability of antihypertensive medicines, team-based care, patient-centred services, and information systems [7].

Early IHCI findings demonstrated that standardized public-sector hypertension management could improve blood



pressure control. Among patients returning for follow-up in sentinel facilities, the proportion with controlled blood pressure increased substantially compared with baseline, with particularly marked improvement in primary-care facilities [16].

Community and primary-care nurses can support such programmes by identifying eligible individuals, measuring blood pressure, maintaining patient records, facilitating follow-up, supporting medication adherence, providing lifestyle counselling, identifying missed appointments, communicating uncontrolled blood pressure to clinicians, and participating in quality-improvement activities.

A 2023 time-motion study examining staff nurses in Indian primary-care facilities highlighted the range of activities involved in hypertension management and provided evidence relevant to workforce planning [17]. The nurse's contribution should therefore be considered when designing staffing patterns and workload allocation for noncommunicable disease services.

Patient-Centred Counselling and Behaviour Change

Successful hypertension control requires patient participation. Community health nurses can promote this by using patient-centred communication rather than directive counselling alone. The nurse can explore what the patient understands about hypertension, what concerns them, what changes they consider achievable, and what support they require.

Motivational communication can be particularly useful when patients are ambivalent about lifestyle change. Small, measurable goals can be established, followed by review during subsequent encounters. For example, instead of simply advising a person to "exercise more," the nurse can collaboratively identify a realistic walking schedule and subsequently review adherence.

Health literacy is another important consideration. Medical terminology should be translated into simple language. Patients should understand why blood pressure needs control, how medicines should be taken, what lifestyle changes are relevant, when follow-up is required, and which warning symptoms require urgent medical attention.

Family and Community Participation

Family involvement can strengthen hypertension management because household practices frequently influence diet, physical activity, medication routines, and healthcare utilization. Nurses can encourage family members to support healthy food preparation, remind patients about appointments, encourage physical activity, and avoid tobacco exposure.

Community-level strategies can extend beyond individual counselling. Nurses can participate in health-awareness campaigns, blood pressure screening days, workplace programmes, community meetings, self-help groups, senior citizen programmes, and local noncommunicable disease initiatives. Community engagement is especially valuable for improving awareness among individuals who may not perceive themselves as being at risk.

Documentation, Monitoring, and Quality Improvement

Documentation is a clinical responsibility as well as a public health function. Community health nurses should accurately record blood pressure readings, screening status, referrals, treatment-related information within their scope, follow-up visits, lifestyle counselling, adherence concerns, and relevant outcomes.

Reliable records allow healthcare teams to identify patients who have missed appointments or remain uncontrolled. At the programme level, aggregated information can reveal gaps in screening coverage, treatment initiation, medicine availability, follow-up, and blood pressure control.

The WHO HEARTS framework includes systems for monitoring because effective hypertension programmes require information for quality improvement rather than simply data collection [9]. Community nurses can therefore contribute directly to population-level quality improvement by ensuring that information generated during routine care is complete and usable.

Digital Health and Future Community Nursing Practice

Digital technology offers new possibilities for hypertension management. Mobile-phone reminders, electronic registers,



teleconsultation, remote blood pressure monitoring, digital education, and automated follow-up alerts can support continuity of care. Nurse-led digital interventions have demonstrated promising effects on blood pressure and self-management outcomes [14].

However, digital health should complement rather than replace human interaction. Older adults, people with limited digital literacy, individuals without smartphones, and economically disadvantaged populations may have difficulty accessing digital services. Community nurses can help reduce this digital divide by providing assistance, combining digital and face-to-face follow-up, and ensuring that technology does not become another barrier to care.

Barriers to Effective Community-Based Hypertension Control

Several challenges may limit the effectiveness of community nursing interventions. These include inadequate staffing, high workload, insufficient training, lack of standardized protocols, poor availability of validated blood pressure devices, medication shortages, inadequate referral systems, fragmented documentation, limited transportation, financial constraints, low health literacy, cultural barriers, and poor continuity of care.

At the patient level, asymptomatic disease presents a unique challenge. Individuals may question why lifelong medication is necessary when they feel healthy. Community nurses must therefore repeatedly explain the preventive purpose of treatment without creating fear.

At the system level, nurses require clear responsibilities and appropriate authority. Task sharing should be accompanied by training, supervision, standardized protocols, referral mechanisms, and quality monitoring. WHO recommends structured approaches to treatment and team-based care as part of scalable hypertension programmes [9,12].

Professional Competencies Required for Community Health Nurses

Effective community hypertension care requires a combination of clinical, educational, communication, organizational, and public health competencies. Nurses

should demonstrate competence in standardized blood pressure measurement, cardiovascular risk assessment, patient education, lifestyle counselling, medication adherence support, recognition of danger signs, referral, documentation, and follow-up.

Communication skills are equally important. The nurse must be able to explain chronic disease concepts in understandable language, establish therapeutic relationships, address misconceptions, respect cultural practices, and communicate effectively with families and multidisciplinary professionals.

Community health nurses also require programme-management competencies. These include maintaining registers, identifying defaulters, monitoring indicators, participating in audits, organizing outreach activities, and using available digital systems. Continuous professional development should incorporate updated hypertension guidelines and locally approved treatment protocols.

Ethical Considerations

Hypertension screening and community-based management must respect autonomy, confidentiality, informed participation, and professional boundaries. Individuals should receive appropriate information about screening and should not be coerced into testing or treatment. Personal health information collected during community visits must be protected.

Nurses should avoid stigmatizing individuals because of obesity, tobacco use, alcohol consumption, dietary practices, or treatment non-adherence. Instead, care should emphasize respect, shared decision-making, and realistic behavioural support. The nurse must also recognize professional limits and avoid making treatment decisions beyond the scope authorized by institutional or national regulations.

Future Directions

Future community hypertension programmes should increasingly combine population screening with systematic follow-up and risk-based management. Training community nurses in standardized blood pressure measurement, cardiovascular risk assessment, behavioural counselling,



digital documentation, and patient-centred communication can strengthen the primary-care workforce.

The expansion of team-based care should also be accompanied by clearly defined roles for nurses, community health workers, physicians, pharmacists, and other healthcare professionals. Standardized protocols can reduce unnecessary variation and facilitate consistent care. WHO's HEARTS approach demonstrates how standardized protocols, essential medicines, team-based care, patient-centred services, and monitoring systems can be integrated into primary healthcare [9].

Research should increasingly examine the long-term sustainability and cost-effectiveness of community nurse-led hypertension programmes. Studies should also consider rural populations, urban poor communities, older adults, people with multiple chronic diseases, and populations experiencing limited healthcare access. Implementation research is particularly important because an intervention that is effective under controlled research conditions may require adaptation before it can be sustained within routine community health services.

Conclusion

Community health nurses have a central role in addressing the hypertension care continuum, beginning with awareness and detection and extending through treatment support, lifestyle modification, adherence, monitoring, referral, and long-term follow-up. Their proximity to individuals and families gives them opportunities to identify undiagnosed hypertension, reinforce healthy behaviours, recognize barriers to treatment, and maintain continuity of care.

The available evidence indicates that structured nurse-led interventions can improve blood pressure outcomes and related self-management behaviours, although effectiveness depends on intervention design, healthcare context, training, professional authority, and integration with the wider healthcare team [3,4]. Community health worker programmes further demonstrate the potential of decentralized screening and follow-up in resource-constrained settings [5,6]. Indian experience through the IHCI provides additional evidence that standardized protocols,

team-based care, patient-centred management, reliable medicines, and information systems can strengthen hypertension control [7,16].

The future of hypertension management therefore lies not only in the availability of effective medicines but also in the development of accessible, continuous, community-oriented systems of care. Community health nurses can serve as an important bridge between individuals, families, communities, and primary healthcare services. By combining accurate assessment with health education, behavioural support, treatment adherence assistance, home-based follow-up, referral, documentation, and digital tools, community health nurses can contribute substantially to earlier detection and sustained control of hypertension and, consequently, to the prevention of cardiovascular and renal complications.

Bibliography

1. World Health Organization. Global report on hypertension: the race against a silent killer. Geneva: World Health Organization; 2023.
2. World Health Organization. First WHO report details devastating impact of hypertension and ways to stop it. Geneva: WHO; 2023.
3. Clark CE, Smith LFP, Taylor RS, Campbell JL. Nurse led interventions to improve control of blood pressure in people with hypertension: systematic review and meta-analysis. *BMJ*. 2010;341:c3995. doi:10.1136/bmj.c3995.
4. Stephen C, Halcomb E, Fernandez R, McInnes S, Batterham M, Zwar N. Nurse-led interventions to manage hypertension in general practice: a systematic review and meta-analysis. *J Adv Nurs*. 2022;78(5):1281-1293. doi:10.1111/jan.15159.
5. Mbuthia GW, Magutah K, Pellowski J. Approaches and outcomes of community health worker's interventions for hypertension management and control in low-income and middle-income countries: systematic review. *BMJ Open*. 2022;12(4):e053455. doi:10.1136/bmjopen-2021-053455.
6. Garg S, Dewangan M, Nanda P, Sahu A, Xalxo L, Bebarta KK, et al. Impact of community health workers on improving identification and primary care of hypertension among the urban poor—findings from Chhattisgarh state of India. *BMC Prim Care*. 2023;24(1):272. doi:10.1186/s12875-023-02231-1.
7. Kaur P, Kunwar A, Sharma M, Durgad K, Gupta S, Bhargava B, et al. The India Hypertension Control



- Initiative—early outcomes in 26 districts across five states of India, 2018-2020. *J Hum Hypertens.* 2023;37(7):560-567. doi:10.1038/s41371-022-00742-5.
8. NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in hypertension prevalence and progress in treatment and control from 1990 to 2019: a pooled analysis of 1201 population-representative studies with 104 million participants. *Lancet.* 2021;398(10304):957-980. doi:10.1016/S0140-6736(21)01330-1.
 9. World Health Organization. HEARTS: Technical package for cardiovascular disease management in primary health care: team-based care. Geneva: World Health Organization; 2018.
 10. World Health Organization. HEARTS: Technical package for cardiovascular disease management in primary health care: healthy-lifestyle counselling. Geneva: World Health Organization; 2018.
 11. Dalal J, Sethi KK, Guha S, Ray S, Deb PK, Kirpalani A, et al. Screening for hypertension in asymptomatic individuals in India: an expert consensus statement. *J Assoc Physicians India.* 2020;68(4):73-79.
 12. World Health Organization. Guideline for the pharmacological treatment of hypertension in adults. Geneva: World Health Organization; 2021.
 13. Zhu X, Wong FKY, Wu CLH. Development and evaluation of a nurse-led hypertension management model in a community: a pilot randomized controlled trial. *Int J Clin Exp Med.* 2015;8(5):? [Community nurse-led intervention study].
 14. The effect of nurse-led digital health interventions on blood pressure control for people with hypertension: a systematic review and meta-analysis. 2023.
 15. World Health Organization. HEARTS: Technical package for cardiovascular disease management in primary health care: risk-based CVD management. Geneva: World Health Organization; 2020.
 16. Chow CK, Teo KK, Rangarajan S, Islam S, Gupta R, Avezum A, et al. Prevalence, awareness, treatment, and control of hypertension in rural and urban communities in low-income and middle-income countries. *JAMA.* 2013;310(9):959-968.
 17. Krishna A, Murali S, Moran AE, Saxena A, Gill SS, Hering D, Kaur P. Understanding the role of staff nurses in hypertension management in primary care facilities in India: a time-motion study. *Prev Chronic Dis.* 2023;20:E39. doi:10.5888/pcd20.220232.
 18. Miao JH, Wang HS, Liu N. The evaluation of a nurse-led hypertension management model in an urban community healthcare: a randomized controlled trial. *Medicine (Baltimore).* 2020;99(27):e20967. doi:10.1097/MD.00000000000020967.
 19. World Health Organization. HEARTS: Technical package for cardiovascular disease management in primary health care: evidence-based treatment protocols. Geneva: World Health Organization; 2018.
 20. World Health Organization. Improving hypertension control in 3 million people: country experiences of programme development and implementation. Geneva: World Health Organization; 2020.
 21. Zhou B, Perel P, Mensah GA, Ezzati M. Global epidemiology, health burden and effective interventions for elevated blood pressure and hypertension. *Nat Rev Cardiol.* 2021;18(11):785-802.
 22. Mills KT, Stefanescu A, He J. The global epidemiology of hypertension. *Nat Rev Nephrol.* 2020;16(4):223-237.
 23. Jeemon P, Séverin T, Amodeo C, Balabanova D, Campbell NRC, Gaita D, et al. World Heart Federation Roadmap for Hypertension—2021 update. *Glob Heart.* 2021;16(1):63. doi:10.5334/gh.1066.
 24. World Health Organization. HEARTS: Technical package for cardiovascular disease management in primary health care: systems for monitoring. Geneva: World Health Organization; 2018.
 25. Kaur P, et al. India Hypertension Control Initiative: hypertension treatment and blood pressure control in a cohort in 24 sentinel site clinics. *J Clin Hypertens.* 2021.