BRIO INNOVATIVE JOURNAL OF NOVEL RESEARCH
BIJNR
Peer Reviewed Indexed Journal
www.bijnr.in
OPEN ACCESS JOURNAL GOOGLE SCHOLAR INDEXED ACADEMIC RESEARCH JOURNAL

www.bijnr.in YEAR: 2024 VOLUME: 1 ISSUE: 2 0766-3242

"Mental Health Nursing in the Era of Rising Adolescent Mental Health Problems"

Karmvir Kaur[1], Dr. Payal Sharma[2]
[1]PhD Research Scholar, [2]Research Supervisor
[1,2]Index Nursing College,
Malwanchal University, Indore

Date of publication: 18/08/2024
DOI 10.5281/zenodo.24467031

Abstract: Adolescent mental health has become an increasingly important public health concern, with psychological, behavioural and emotional difficulties affecting the wellbeing, academic performance, interpersonal relationships and future development of young people. Adolescence represents a critical developmental period characterised by substantial biological, psychological and social transitions. Exposure to academic pressure, family conflicts, peer victimisation, cyberbullying, substance use, social media influences, socioeconomic difficulties and traumatic experiences may increase vulnerability to mental health problems. Depression, anxiety disorders, behavioural disorders, eating disorders, substance-related problems, self-harm and suicidal behaviour are among the major concerns affecting this population. Mental health nursing has an essential role in addressing these challenges through health promotion, early identification, comprehensive assessment, therapeutic communication, crisis intervention, family counselling and rehabilitation. This narrative review examines the changing landscape of adolescent mental health problems and explores the implications for contemporary mental health nursing practice. Particular attention is given to developmental considerations, emerging psychosocial risk factors, common psychiatric conditions, nursing assessment, preventive interventions, school-based mental health services, family-centred care and multidisciplinary collaboration. The review also highlights the importance of mental health literacy, digital mental health support, suicide prevention and community-based interventions. Mental health nurses are strategically positioned to bridge the gap between adolescents experiencing psychological difficulties and accessible mental healthcare services. Strengthening nursing competencies, integrating mental health services into educational and community settings, reducing stigma and promoting early intervention are essential for improving adolescent mental health outcomes. A comprehensive, developmentally appropriate and evidence-informed nursing approach can contribute to early recovery, psychological resilience and healthier transitions into adulthood.

Keywords: *Adolescent mental health, mental health nursing, depression, anxiety, psychological wellbeing, psychiatric nursing, suicide prevention, mental health promotion, early intervention, school mental health, psychosocial assessment, adolescent counselling, resilience, community mental health.*

1. Introduction

Adolescence is a significant developmental stage involving physical maturation, cognitive development, emotional changes and the formation of social identity. During this period, individuals gradually develop independence, establish interpersonal relationships and acquire skills necessary for adulthood. Although adolescence offers considerable opportunities for growth, it is also a period of increased vulnerability to psychological and behavioural difficulties. Mental health problems emerging during this stage may interfere with educational achievement, social functioning, family relationships and long-term quality of life.

The global burden of adolescent mental health problems has received increasing attention from healthcare professionals and policymakers. According to the World Health Organization and UNICEF estimates published before 2024, approximately one in seven adolescents aged 10–19 years experiences a mental disorder [1,2]. Depression, anxiety and behavioural disorders contribute substantially to the burden of illness and disability in this age group. Many mental health conditions begin during adolescence or early adulthood,

BRIO INNOVATIVE JOURNAL OF NOVEL RESEARCH
BIJNR
Peer Reviewed Indexed Journal
www.bijnr.in
OPEN ACCESS JOURNAL GOOGLE SCHOLAR INDEXED ACADEMIC RESEARCH JOURNAL
www.bijnr.in YEAR: 2024 VOLUME: 1 ISSUE: 2 0766-3242

making early recognition and intervention particularly important [3].

The increasing complexity of adolescent lifestyles has introduced additional challenges to psychological wellbeing. Academic competition, examination anxiety, unemployment-related family stress, changing family structures, social media exposure, cyberbullying, body image dissatisfaction and reduced opportunities for meaningful social interaction may contribute to emotional distress. The COVID-19 pandemic further highlighted the importance of adolescent mental health by disrupting education, social relationships, recreational activities and access to support services.

Mental health nursing has traditionally focused on the assessment and management of individuals experiencing psychiatric disorders. However, contemporary adolescent mental health nursing requires a broader approach that includes prevention, psychological wellbeing, developmental support, early identification, family education and community participation. Nurses working in psychiatric hospitals, general hospitals, schools, primary healthcare facilities and community settings have important opportunities to identify psychological difficulties before they progress into severe mental health conditions.

2. Emerging Landscape of Adolescent Mental Health Problems

The pattern of adolescent mental health problems is influenced by interactions among biological, psychological, family-related, educational, cultural and environmental factors. Anxiety and depressive disorders are major concerns, while behavioural disorders, attention-deficit/hyperactivity disorder, eating disorders, substance-related problems and self-harming behaviours also contribute to adolescent morbidity [1,4].

Depression in adolescents may present through persistent sadness, irritability, loss of interest, reduced motivation, feelings of worthlessness, sleep disturbances and declining academic performance. Unlike some adult presentations, adolescents may demonstrate irritability, social withdrawal or behavioural changes rather than clearly communicating feelings of sadness. Anxiety disorders may manifest through excessive worrying, fear of failure, avoidance of social situations, panic symptoms and physical complaints such as headaches or abdominal discomfort.

Behavioural and emotional difficulties may include aggression, impulsivity, oppositional behaviour, conduct-related problems and difficulties regulating emotions. Attention-deficit/hyperactivity disorder may affect concentration, academic performance, organisation and interpersonal relationships. Eating disorders and body image disturbances are additional concerns, particularly among adolescents exposed to unrealistic appearance standards and excessive social comparison.

Substance use represents another important challenge. Experimentation with alcohol, tobacco, nicotine products and other psychoactive substances may occur during adolescence and can contribute to behavioural, psychological and social complications. Mental health nurses should assess substance use without judgement and consider its relationship with underlying emotional difficulties.

The Global Burden of Disease research has demonstrated the substantial contribution of mental disorders to disability among children, adolescents and young adults [4]. These findings reinforce the need for adolescent mental health services that integrate prevention, early detection and continuing care.

3. Factors Contributing to Rising Adolescent Mental Health Problems

Adolescent psychological wellbeing is shaped by multiple interacting determinants. Understanding these factors enables mental health nurses to conduct comprehensive assessments and develop interventions that address the underlying causes of distress rather than focusing exclusively on symptoms.

Academic and educational pressures: Academic competition, examination-related anxiety, parental expectations, fear of failure and uncertainty regarding future careers can produce significant psychological stress. Adolescents may experience sleep disturbances, reduced self-confidence and emotional exhaustion when academic achievement becomes the primary measure of personal success. Nurses should assess academic stress as part of routine psychosocial evaluation.

BRIO INNOVATIVE JOURNAL OF NOVEL RESEARCH
BIJNR
Peer Reviewed Indexed Journal
www.bijnr.in
OPEN ACCESS JOURNAL GOOGLE SCHOLAR INDEXED ACADEMIC RESEARCH JOURNAL
www.bijnr.in YEAR: 2024 VOLUME: 1 ISSUE: 2 0766-3242

Family-related factors: Family conflict, parental separation, domestic violence, neglect, inadequate emotional support and unrealistic expectations may negatively affect adolescent mental health. Supportive family relationships, effective communication and positive parenting practices are important protective factors. Family assessment should therefore be incorporated into adolescent mental health nursing.

Digital technology and social media: Digital platforms influence adolescent communication, identity formation and social relationships. Excessive screen use, cyberbullying, online harassment, exposure to harmful content and unrealistic social comparisons may contribute to anxiety, sleep disturbances and low self-esteem. However, digital platforms can also provide opportunities for social support and mental health education. Nurses should promote balanced and responsible technology use rather than assuming that all digital engagement is harmful.

Peer relationships and social exclusion: Peer acceptance becomes increasingly important during adolescence. Bullying, discrimination, friendship conflicts, romantic relationship difficulties and social isolation may contribute to emotional distress. Adolescents who experience repeated rejection may develop negative self-perceptions and avoid social interactions.

Socioeconomic and environmental influences: Poverty, food insecurity, inadequate housing, community violence, limited educational opportunities and restricted access to healthcare may increase psychological vulnerability. The Lancet Commission on adolescent health emphasised the importance of addressing the broader social and environmental determinants of adolescent wellbeing [3].

Biological and developmental factors: Hormonal changes, neurological development, genetic vulnerability, sleep patterns and individual temperament influence adolescent emotional regulation and psychological functioning. Nurses should recognise that behavioural changes may reflect developmental or psychiatric difficulties requiring further assessment.

4. Implications for Mental Health Nursing Assessment

Comprehensive assessment is the foundation of effective adolescent mental health nursing. Adolescents may express psychological distress through behavioural changes, physical complaints, academic difficulties or social withdrawal rather than directly reporting emotional symptoms. Therefore, nurses should use a developmentally appropriate and holistic assessment approach.

Establishing therapeutic relationships: Establishing trust is essential before discussing sensitive psychological concerns. Nurses should provide a private and supportive environment, demonstrate empathy, maintain non-judgmental communication and respect the adolescent's experiences. Open-ended questions encourage adolescents to express their feelings and concerns without fear of criticism.

Psychosocial assessment: Assessment should explore family relationships, peer interactions, academic performance, social participation, lifestyle habits, traumatic experiences and available support systems. Nurses should also consider cultural beliefs, socioeconomic circumstances and environmental stressors that may influence mental health.

Mental status examination: A systematic mental status examination should assess appearance, behaviour, psychomotor activity, speech, mood, affect, thought processes, thought content, perception, cognition, insight and judgement. Developmental considerations should be incorporated when interpreting findings.

Identification of psychological symptoms: Nurses should assess depressive symptoms, anxiety, irritability, sleep disturbances, appetite changes, concentration difficulties, hopelessness and loss of interest. Validated instruments such as the Patient Health Questionnaire-9 modified for adolescents (PHQ-A), Generalized Anxiety Disorder-7 (GAD-7) and Strengths and Difficulties Questionnaire (SDQ) may support assessment when appropriately used. Screening results should not replace clinical evaluation.

Suicide and self-harm assessment: Direct assessment of suicidal thoughts, previous attempts, self-harming behaviours, intent, planning and access to potentially harmful means is essential when indicated. The WHO mhGAP

BRIO INNOVATIVE JOURNAL OF NOVEL RESEARCH
BIJNR
Peer Reviewed Indexed Journal
www.bijnr.in
OPEN ACCESS JOURNAL GOOGLE SCHOLAR INDEXED ACADEMIC RESEARCH JOURNAL
www.bijnr.in YEAR: 2024 VOLUME: 1 ISSUE: 2 0766-3242

Intervention Guide recommends assessment of self-harm and suicidal thoughts in relevant high-risk clinical situations [5]. Adolescents with immediate safety concerns require urgent clinical evaluation, appropriate supervision and activation of emergency protocols.

Assessment of protective factors: Nurses should identify positive family relationships, supportive friendships, personal strengths, coping abilities, cultural resources, educational support and future aspirations. Recognising protective factors helps develop individualised interventions that strengthen resilience.

5. Mental Health Nursing Interventions for Adolescents

Mental health nursing interventions should address immediate psychological difficulties while promoting long-term emotional development and functional recovery. The approach should be individualised according to developmental stage, psychiatric symptoms, family circumstances and available resources.

5.1 Mental Health Promotion and Primary Prevention

Primary prevention aims to strengthen psychological wellbeing before significant mental health problems develop. Nurses can organise educational sessions on emotional regulation, stress management, healthy relationships, problem-solving, sleep hygiene and responsible digital behaviour.

Life-skills education is particularly valuable because it enables adolescents to develop practical abilities for managing everyday challenges. WHO's *Helping Adolescents Thrive* guidelines recommend evidence-informed psychosocial interventions to promote mental health, prevent mental disorders and reduce self-harm and other risk behaviours [6]. Nurses can contribute to implementing these approaches through schools, community programmes and healthcare services.

5.2 Early Identification and Secondary Prevention

Secondary prevention focuses on identifying emerging mental health difficulties and initiating timely interventions. Nurses should recognise warning signs such as persistent sadness, social withdrawal, unexplained behavioural changes, declining academic performance, repeated physical complaints and sleep disturbances.

Routine adolescent health consultations provide opportunities to identify psychological concerns. Early referral to mental health professionals can prevent unnecessary delays in treatment. Community and school health nurses should establish clear referral pathways for adolescents requiring specialised assessment.

5.3 Therapeutic Communication and Supportive Counselling

Therapeutic communication helps adolescents express emotional difficulties and develop a sense of psychological security. Active listening, clarification, reflection, reassurance within appropriate limits and emotional validation are useful nursing techniques.

Supportive counselling can address examination anxiety, relationship difficulties, family conflicts and adjustment problems. Nurses should avoid minimising adolescent concerns or comparing their experiences with those of others. Instead, they should acknowledge individual experiences and support the development of realistic coping strategies.

5.4 Stress Management and Resilience Development

Resilience refers to the ability to adapt and recover when confronted with adversity. Nurses can encourage relaxation exercises, mindfulness-based activities, structured physical activity, creative expression, positive self-talk and problem-solving techniques.

Adolescents should be supported in identifying personal strengths, setting achievable goals and maintaining healthy relationships. Resilience programmes should not imply that individuals alone are responsible for overcoming adversity; supportive families, safe schools and accessible health services are equally important.

5.5 Management of Depression and Anxiety

Adolescents experiencing depression or anxiety may require psychological treatment, medication or combined interventions depending on clinical severity and professional assessment. Mental health nurses can monitor symptoms, provide psychoeducation, support treatment adherence and identify adverse effects of prescribed medication.

Nurses can also reinforce psychological treatment strategies and encourage participation in appropriate counselling.

BRIO INNOVATIVE JOURNAL OF NOVEL RESEARCH
BIJNR
Peer Reviewed Indexed Journal
www.bijnr.in
OPEN ACCESS JOURNAL GOOGLE SCHOLAR INDEXED ACADEMIC RESEARCH JOURNAL
www.bijnr.in YEAR: 2024 VOLUME: 1 ISSUE: 2 0766-3242

Severe symptoms, psychotic features, significant functional impairment or suicidal risk require specialist psychiatric assessment.

5.6 Crisis Intervention and Suicide Prevention

Crisis intervention is an essential responsibility of psychiatric and mental health nurses. Adolescents experiencing suicidal thoughts, severe emotional distress, self-harm or acute behavioural disturbances require prompt assessment and safety-focused care.

Nurses should maintain a supportive environment, undertake appropriate risk assessment, involve the relevant mental health team and facilitate emergency referral when required. Safety planning, family involvement when appropriate, restriction of access to potentially harmful means and continuity of follow-up are important components of suicide prevention.

6. Role of School Mental Health Nursing

Schools provide valuable opportunities for promoting psychological wellbeing because they enable regular contact with large adolescent populations. School health nurses can identify early warning signs, provide health education, support students experiencing emotional difficulties and facilitate referral to specialised services.

School-based mental health programmes can address examination anxiety, bullying, cyberbullying, substance use, emotional regulation and interpersonal relationships. Nurses can organise awareness campaigns, counselling activities, peer-support programmes and teacher education sessions.

An integrative review by Hoskote and colleagues highlighted the evolving role of school nurses in addressing adolescent mental health at individual, community and systems levels [7]. This supports the importance of positioning school nurses as contributors to comprehensive student mental health services rather than limiting their responsibilities to physical health concerns.

School mental health programmes should promote psychological safety, confidentiality, inclusion and non-discrimination. Teachers, counsellors, parents and healthcare professionals should collaborate to identify students who require additional support.

7. Family-Centred Mental Health Nursing

Family participation is an important component of adolescent mental healthcare. Parents and caregivers influence emotional development, behaviour, treatment adherence and access to professional support. However, family conflict, criticism and lack of mental health awareness may prevent adolescents from seeking assistance.

Nurses can educate families about common adolescent mental health problems, early warning signs, treatment options and supportive communication. Parents should be encouraged to listen without judgement, avoid excessive criticism and recognise the importance of emotional validation.

Family counselling may help address communication difficulties, interpersonal conflicts and behavioural concerns. Nurses should also recognise that family involvement must be balanced with adolescent privacy, confidentiality, developmental autonomy and safeguarding responsibilities.

Supportive parenting programmes and caregiver education can strengthen family relationships and improve the emotional environment in which adolescents develop. The WHO and UNICEF *Helping Adolescents Thrive Toolkit* emphasises supportive environments and caregiver involvement as important components of adolescent mental health promotion [8].

8. Community Mental Health Nursing and Multidisciplinary Collaboration

Community mental health nursing provides an important connection between adolescents, families, educational institutions and healthcare services. Nurses can conduct community awareness programmes, organise adolescent health education, identify vulnerable individuals and facilitate access to counselling and psychiatric services.

Community-based interventions should address mental health stigma, social exclusion, substance use, bullying and limited access to psychological support. Youth clubs, community centres and educational institutions can serve as platforms for mental health promotion.

Multidisciplinary collaboration is essential because adolescent mental health problems frequently involve medical, psychological, educational and social dimensions. Psychiatrists, psychologists, paediatricians, social workers,

BRIO INNOVATIVE JOURNAL OF NOVEL RESEARCH
BIJNR
Peer Reviewed Indexed Journal
www.bijnr.in
OPEN ACCESS JOURNAL GOOGLE SCHOLAR INDEXED ACADEMIC RESEARCH JOURNAL
www.bijnr.in YEAR: 2024 VOLUME: 1 ISSUE: 2 0766-3242

teachers, counsellors and community organisations should work together to provide coordinated care.

Mental health nurses can act as care coordinators by arranging referrals, monitoring progress, supporting treatment adherence and facilitating communication between healthcare providers and families. The WHO mhGAP framework provides guidance for integrating mental health assessment and management into non-specialised healthcare settings [5].

9. Digital Mental Health and Emerging Nursing Responsibilities

Digital health technologies offer new opportunities for adolescent mental health promotion and service delivery. Teleconsultation, mobile health applications, online counselling, digital educational resources and virtual support programmes may improve access to information and professional assistance.

Mental health nurses can use digital platforms to provide psychoeducation, reinforce coping strategies, support follow-up and communicate with adolescents through appropriately governed healthcare systems. However, digital interventions must consider confidentiality, data protection, informed consent, cyberbullying and the possibility of inappropriate online interactions.

Digital mental health services should complement rather than completely replace face-to-face assessment, particularly when adolescents demonstrate severe psychological symptoms or suicide risk. Nurses should also be equipped to identify misinformation, harmful online content and inappropriate digital health practices.

10. Challenges in Contemporary Adolescent Mental Health Nursing

Several challenges affect the delivery of effective adolescent mental health services. Mental health stigma remains a major barrier, particularly in communities where psychiatric difficulties are associated with shame or misconceptions. Adolescents may conceal symptoms because they fear discrimination, family criticism or social exclusion.

Limited availability of trained mental health professionals, inadequate school counselling facilities, insufficient resources and delayed referral pathways further restrict access to care. In some healthcare settings, mental health assessment may receive less attention than physical health assessment.

Another challenge is the shortage of specialised nursing training in adolescent psychiatric care. Nurses require continuing education in developmental psychology, mental status examination, suicide assessment, therapeutic communication, psychopharmacology, safeguarding and evidence-based interventions.

Cultural sensitivity is equally important. Mental health symptoms may be expressed differently across communities, and beliefs regarding emotional distress and psychiatric treatment can influence help-seeking behaviour. Nursing interventions should therefore be culturally appropriate and responsive to local needs.

11. Future Directions for Mental Health Nursing

The increasing recognition of adolescent mental health as a public health priority requires the development of stronger nursing services and innovative preventive approaches. Mental health nursing education should incorporate specialised adolescent assessment, developmental psychopathology, digital mental health, suicide prevention and family-centred interventions.

Greater integration of mental health services into primary healthcare, schools and community programmes is necessary to improve early detection and reduce treatment gaps. Nurses should participate in developing institutional adolescent mental health policies, referral protocols and preventive programmes.

Nursing research should focus on evaluating school-based interventions, nurse-led counselling, digital mental health programmes, family education and community resilience initiatives. Research from India and other low- and middle-income countries is particularly important for understanding cultural influences, healthcare accessibility and resource-related barriers.

Future nursing practice should also emphasise measurable outcomes, including improved mental health literacy, early identification, reduced psychological distress, treatment adherence, social functioning and quality of life. Such an approach can strengthen the contribution of nursing to

BRIO INNOVATIVE JOURNAL OF NOVEL RESEARCH
BIJNR
Peer Reviewed Indexed Journal
www.bijnr.in
OPEN ACCESS JOURNAL GOOGLE SCHOLAR INDEXED ACADEMIC RESEARCH JOURNAL
www.bijnr.in YEAR: 2024 VOLUME: 1 ISSUE: 2 0766-3242

adolescent mental health at individual, institutional and population levels.

12. Conclusion

The increasing burden of adolescent mental health problems represents a significant challenge for contemporary healthcare systems. Depression, anxiety, behavioural disturbances, substance use, self-harm and other psychological difficulties can interfere with adolescent development and produce consequences that extend into adulthood. Biological changes, academic pressures, family relationships, peer influences, digital environments and socioeconomic circumstances collectively shape adolescent mental health.

Mental health nursing has an essential role in responding to these challenges through health promotion, comprehensive assessment, early identification, therapeutic communication, crisis intervention, family counselling and rehabilitation. School health nurses and community mental health nurses are particularly important in extending preventive services beyond psychiatric hospitals and specialised healthcare facilities.

An effective response requires a shift from a predominantly treatment-oriented model towards a comprehensive approach integrating promotion, prevention, early intervention and continuing care. Strengthening nursing education, developing adolescent-friendly services, reducing stigma and improving multidisciplinary collaboration are necessary steps towards achieving this objective.

Ultimately, mental health nursing should aim not only to manage psychiatric symptoms but also to empower adolescents to develop emotional resilience, healthy relationships, self-confidence and effective coping abilities. By recognising adolescents as active participants in their own healthcare and creating supportive environments across families, schools and communities, mental health nurses can contribute substantially to healthier transitions into adulthood.

References

1. World Health Organization. Adolescent mental health. Geneva: World Health Organization; 2021.
2. UNICEF. The State of the World's Children 2021: On My Mind – Promoting, Protecting and Caring for Children's Mental Health. New York: United Nations Children's Fund; 2021.
3. Patton GC, Sawyer SM, Santelli JS, Ross DA, Afifi R, Allen NB, et al. Our future: a Lancet commission on adolescent health and wellbeing. Lancet. 2016;387(10036):2423-2478. doi:10.1016/S0140-6736(16)00579-1.
4. Erskine HE, Sweeney GM, O'Connell CM, et al. Worldwide prevalence and disability from mental disorders across childhood and adolescence: evidence from the Global Burden of Disease Study. JAMA Psychiatry. 2024.
5. World Health Organization. mhGAP Intervention Guide for Mental, Neurological and Substance Use Disorders in Non-Specialized Health Settings. Version 2.0. Geneva: WHO; 2016.
6. World Health Organization. Guidelines on Mental Health Promotive and Preventive Interventions for Adolescents: Helping Adolescents Thrive. Geneva: WHO; 2020.
7. Hoskote AR, Croce E, Johnson KE. The evolution of the role of U.S. school nurses in adolescent mental health at the individual, community, and systems level: an integrative review. J Sch Nurs. 2023;39(1):51-71. doi:10.1177/10598405211068120.
8. World Health Organization, United Nations Children's Fund. Helping Adolescents Thrive Toolkit: Strategies to Promote and Protect Adolescent Mental Health and Reduce Self-Harm and Other Risk Behaviours. Geneva: WHO and UNICEF; 2021.
9. Kieling C, Baker-Henningham H, Belfer M, Conti G, Ertem I, Omigbodun O, et al. Child and adolescent mental health worldwide: evidence for action. Lancet. 2011;378(9801):1515-1525. doi:10.1016/S0140-6736(11)60827-1.
10. Gore FM, Bloem PJN, Patton GC, Ferguson J, Joseph V, Coffey C, et al. Global burden of disease in young people aged 10-24 years: a systematic analysis. Lancet. 2011;377(9783):2093-2102. doi:10.1016/S0140-6736(11)60512-6.
11. Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity

BRIO INNOVATIVE JOURNAL OF NOVEL RESEARCH
BIJNR
Peer Reviewed Indexed Journal
www.bijnr.in
OPEN ACCESS JOURNAL GOOGLE SCHOLAR INDEXED ACADEMIC RESEARCH JOURNAL
www.bijnr.in YEAR: 2024 VOLUME: 1 ISSUE: 2 0766-3242

Survey Replication. Arch Gen Psychiatry. 2005;62(6):593-602. doi:10.1001/archpsyc.62.6.593.
12. Merikangas KR, He JP, Burstein M, Swanson SA, Avenevoli S, Cui L, et al. Lifetime prevalence of mental disorders in U.S. adolescents: results from the National Comorbidity Survey Replication-Adolescent Supplement. J Am Acad Child Adolesc Psychiatry. 2010;49(10):980-989. doi:10.1016/j.jaac.2010.05.017.
13. Solmi M, Radua J, Olivola M, Croce E, Soardo L, Salazar de Pablo G, et al. Age at onset of mental disorders worldwide: large-scale meta-analysis of 192 epidemiological studies. Mol Psychiatry. 2022;27:281-295. doi:10.1038/s41380-021-01161-7.
14. World Health Organization. Health for the World's Adolescents: A Second Chance in the Second Decade. Geneva: WHO; 2014.
15. World Health Organization. Preventing Suicide: A Global Imperative. Geneva: WHO; 2014.
16. World Health Organization. Comprehensive Mental Health Action Plan 2013–2020. Geneva: WHO; 2013.
17. World Health Organization. Global Accelerated Action for the Health of Adolescents (AA-HA!): Guidance to Support Country Implementation. Geneva: WHO; 2017.
18. World Health Organization. Guidelines on Mental Health Promotive and Preventive Interventions for Adolescents: Helping Adolescents Thrive – Executive Summary. Geneva: WHO; 2021.
19. National Academies of Sciences, Engineering, and Medicine. The Promise of Adolescence: Realizing Opportunity for All Youth. Washington, DC: National Academies Press; 2019.
20. Sawyer SM, Afifi RA, Bearinger LH, Blakemore SJ, Dick B, Ezeh AC, et al. Adolescence: a foundation for future health. Lancet. 2012;379(9826):1630-1640. doi:10.1016/S0140-6736(12)60072-5.
21. American Academy of Pediatrics. Promoting optimal development: identifying infants and young children with developmental disorders through developmental surveillance and screening. Pediatrics. 2020;145(1):e20193449.
22. World Health Organization. INSPIRE: Seven Strategies for Ending Violence Against Children. Geneva: WHO; 2016.
23. World Health Organization. Guidelines for the Management of Physical Health Conditions in Adults with Severe Mental Disorders. Geneva: WHO; 2018.
24. UNICEF. The State of the World's Children 2021: On My Mind – Summary Report. New York: United Nations Children's Fund; 2021.
25. Patel V, Flisher AJ, Hetrick S, McGorry P. Mental health of young people: a global public-health challenge. Lancet. 2007;369(9569):1302-1313. doi:10.1016/S0140-6736(07)60368-7.
26. World Health Organization. Preventing Suicide: A Resource for Teachers and Other School Staff. Geneva: WHO; 2000.
27. World Health Organization. Mental Health Action Plan 2013–2030. Geneva: WHO; 2021.